

Surgical Pathology Report

SPECIMEN(S) RECEIVED

- A: Cervical biopsy 5 o'clock
- B: Left ovarian lesion
- C: Left ovarian fossa, biopsy
- D: Left periureteral, biopsy
- E: Left uterosacral ligament, biopsy
- F: Left perirectal, biopsy
- G: Left cul de sac, biopsy
- H: Midline cul de sac, biopsy
- I: Right cul de sac, biopsy
- J: Right perirectal, biopsy
- K: Right periureteral, biopsy

CLINICAL INFORMATION

Abnormal estrogen symptoms postprocedural ovarian failure

PRE OP DIAGNOSIS

ABNORMAL ESTROGEN SYMPTOMS

FINAL DIAGNOSIS

- A. Cervical 5:00, biopsy:
 - Endocervical tissue within normal limits.
 - Very scant squamous epithelium (0.1 mm in size), negative for high-grade dysplasia.
 - See comment.

COMMENT:

There is a very short segment of squamous epithelium, negative for p16. It most likely represents basal layer of normal squamous epithelium, negative for high-grade dysplasia.

- B. Left ovarian lesion, biopsy:
 - Ovarian tissue with feature of endometrioma.
- C. Left ovarian fossa, biopsy:
 - Involved by endometriosis.
- D. Left periureteral, biopsy:
 - Involved by endometriosis.
- E. Left uterosacral ligament, biopsy:
 - Fibroadipose tissue within normal limits.
- F. Left perirectal, biopsy:
 - Fibroadipose tissue with granulomatous inflammation and multinucleated giant cells.

- G. Left cul de sac, biopsy:
Involved by endometriosis.
- H. Midline cul de sac, biopsy:
Fibroadipose tissue with very focal feature of possible endometriosis.
- I. Right cul de sac, biopsy:
Involved by endometriosis.
- J. Right perirectal, biopsy:
Involved by endometriosis.
- K. Right periureteral, biopsy:
Fibroadipose tissue within normal limits.

GROSS DESCRIPTION

The specimen is received in eleven parts.

Part A, received in formalin is (are) 2 piece(s) of tan white soft tissue measuring 0.4 cm. in greatest dimension. Entirely submitted in A1.

Part B, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 0.7 cm. in greatest dimension. Entirely submitted in B1.

Part C, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 0.7 cm. in greatest dimension. Entirely submitted in C1.


Part D, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 1.0 cm. in greatest dimension. Entirely submitted in D1.

Part E, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 1.0 cm. in greatest dimension. Entirely submitted in E1.

Part F, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 0.8 cm. in greatest dimension. Entirely submitted in F1.

Part G, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 2.0 cm. in greatest dimension. Entirely submitted in G1.

Part H, received in formalin is (are) 1 piece(s) of tan yellow soft tissue measuring 1.0 cm. in greatest dimension. Entirely submitted in H1.



Part I, received in formalin is (are) 1 piece(s) of pale tan soft tissue measuring 1.0 cm. in greatest dimension. Entirely submitted in I1.

Part J, received in formalin is (are) 1 piece(s) of yellow soft tissue measuring 0.6 cm. in greatest dimension. Entirely submitted in J1.

Part K, received in formalin is (are) 1 piece(s) of tan red soft tissue measuring 1.6 cm. in greatest dimension. Entirely submitted in K1.



MICROSCOPIC DESCRIPTION

Unless "gross only" is specified, the final diagnosis for each specimen is based on a microscopic examination of representative sections of tissue.

