

Histology Cytology Report

Clinical Information

Preoperative diagnosis: Adenomyosis

Clinical data:

Special instructions:

Specimen source: A. Uterus, bilateral fallopian tubes; B. Right cul-de-sac

Diagnosis

A. UTERUS AND BILATERAL FALLOPIAN TUBES, EXCISION:

INACTIVE ENDOMETRIUM.

FOCAL SUPERFICIAL ADENOMYOSIS UTERI.

UTERINE SEROSAL ADHESIONS.

BILATERAL FALLOPIAN TUBES WITH NO SIGNIFICANT HISTOPATHOLOGIC ABNORMALITY.

B. RIGHT PELVIC CUL-DE-SAC, EXCISION:

SUBPERITONEAL FIBROSIS.

Gross Description

A. In formalin labeled "bilateral fallopian tubes, uterus" is a supracervical uterus and detached bilateral fallopian tubes. The uterus is 47 g, 5.0 cm endocervix to fundus, 5.5 cm cornu to cornu and 3 cm anterior to posterior. The serosal surface is tan-gray with focal hemorrhagic adhesions on the anterior surface. The uterine cavity is a 4.0 x 2.5 cm. The tan-red endometrium is 0.1 cm thick and overlies tan-pink myometrium that is congested towards the serosa. The fimbriated fallopian tubes are 6 cm length by 0.5 cm diameter. Representative sections submitted as follows: 1-anterior endocervical margin, 2-posterior endocervical margin, 3-anterior endomyometrium, 4-posterior endomyometrium, 5-6 - representative from both fallopian tubes including entire fimbriated ends, 7-anterior serosa with hemorrhagic adhesions.

B. In formalin labeled "right cul-de-sac" are 3 tan soft tissue fragments from 0.8 x 0.7 x 0.3 cm to 1.0 x 0.5 x 0.3 cm. Totally submitted, one block. JHA

Microscopic Description

A and B. Gross description reviewed and microscopic examined.