

Clinical Information

Preoperative diagnosis: Endometriosis and uterine fibroids

Clinical data:

Postoperative diagnosis: Pending

Special instructions:

Specimen source: A. Left ovarian fossa; B. Left periureteral; C. Left uterine artery; D. Left uterosacral ligament; E. Left rectal wall; F. Right ovarian fossa; G. Right periureteral; H. Uterine fibroids

Diagnosis

A. LEFT OVARIAN FOSSA, EXCISION:

MILD SUBPERITONEAL FIBROSIS AND ECTOPIC DECIDUAL REACTION.

- B. LEFT PERIURETERAL PERITONEUM, EXCISION: ENDOMETRIOSIS WITH ECTOPIC DECIDUAL REACTION.
- C. LEFT UTERINE ARTERY, EXCISION: ENDOMETRIOSIS WITH ECTOPIC DECIDUAL REACTION.
- D. LEFT UTEROSACRAL LIGAMENT, EXCISION: ENDOMETRIOSIS WITH ECTOPIC DECIDUAL REACTION.
- E. LEFT RECTAL WALL, EXCISION: ECTOPIC DECIDUAL REACTION.
- F. RIGHT OVARIAN FOSSA, EXCISION:
 MILD SUBPERITONEAL FIBROSIS AND FOCAL ECTOPIC DECIDUAL REACTION.
- G. RIGHT PERIURETERAL PERITONEUM: ECTOPIC DECIDUAL REACTION.
- H. UTERINE FIBROIDS, EXCISION:
 PARTIALLY INFARCTED LEIOMYOMA(S).

Diagnosis

Gross Description

- A. In formalin labeled "left ovarian fossa": One, 1 x 0.7×0.3 cm tissue piece(s). Totally submitted, 1 block.
- B. In formalin labeled "left periureteral": One, $1.8 \times 0.7 \times 0.5$ cm tissue piece(s). Specimen is sectioned. Totally submitted, 1 block.
- C. In formalin labeled "left uterine artery": One, $0.9 \times 0.5 \times 0$
- D. In formalin labeled "left uterosacral ligament": One, $2.2 \times 0.7 \times 0.3$ cm tissue piece(s). Specimen is sectioned. Totally submitted, 1 block.
- E. In formalin labeled "left rectal wall": One, 1.4 \times 0.8 \times 0.4 cm tissue piece(s). Specimen is sectioned. Totally submitted, 1 block.
- F. In formalin labeled "right ovarian fossa": One, $0.7 \times 0.4 \times 0.3$ cm tissue piece(s). Totally submitted, 1 block.
- G. In formalin labeled "right periureteral": One, $1.3 \times 0.6 \times 0.3$ cm tissue piece(s). Specimen is sectioned. Totally submitted, 1 block.
- H. Received in formalin labeled "uterine fibroids" are multiple tan-white, fragmented portions of mass that are 23 g, $7 \times 7 \times 2.5$ cm aggregate. The largest fragment is $5 \times 3 \times 2.5$ cm. Sectioning of the largest fragment reveals white, rubbery, whorled cut surfaces remarkable for focally red, and degenerative areas. Sampled, 4 blocks.

Microscopic Description

A through H. Gross description reviewed and microscopic examined.