

Histology Cytology Report

Preoperative diagnosis: Superficial endometriosis of the anterior cul-de-sac

Clinical data:

Postoperative diagnosis: Pending

Special instructions:

Specimen source: A. Omentum; B. Uterus, cervix, right fallopian tube; C. Left fallopian tube and ovary; D. Left periureteral; E. Right cul-de-sac

### Diagnosis

A. OMENTUM, EXCISION:

FOCAL SURFACE PERITONEAL ADHESIONS.

B. UTERUS, CERVIX, RIGHT FALLOPIAN TUBE, EXCISION:

UTERINE CERVIX WITH NO SIGNIFICANT HISTOPATHOLOGIC ABNORMALITY.

PROLIFERATIVE ENDOMETRIUM.

FOCAL SUPERFICIAL ADENOMYOSIS UTERI.

UTERINE SEROSAL ADHESIONS.

CYSTIC WALTHARD RESTS, FALLOPIAN TUBE.

C. LEFT FALLOPIAN TUBE AND OVARY, EXCISION:

TUBO-OVARIAN ADHESIONS.

BENIGN SEROUS CYSTADENOFIBROMA AND LUTEINIZED FOLLICULAR CYST, OVARY.

D. LEFT PERIURETHRAL PERITONEUM, EXCISION:

**ENDOSALPINGOSIS.** 

E. RIGHT PELVIC CUL-DE-SAC, EXCISION:

ENDOSALPINGOSIS.

#### **Gross Description**

A. In formalin labeled "omentum" is a  $7.5 \times 5 \times 2$  cm portion of finally lobulated omentum that is remarkable for scant adhesions and congestion. Sampled, 2 blocks.



## Histology Cytology Report

## **Gross Description**

- B. In formalin labeled "uterus, cervix, right fallopian tube" is a uterus with amputated fallopian tube. The 109 g uterus is 12 cm fundus to ectocervix, 7 cm cornu to cornu and 5 cm anterior to posterior. The serosal surface is pink-tan with focal congestion and marked adhesion, anteriorly. There is a 3.5 cm diameter cervix surfaced by smooth ectocervical mucosa. Sectioning demonstrates a triangular uterine cavity that is 5 x 2.5 cm and lined by pink-tan, lush, 0.8 cm thick endometrium. The myometrium is extensively trabeculated and 2 cm thick. No discrete masses are identified. The fimbriated fallopian tube is 7 x 0.5 cm with congested serosa. Sampled: 1-2-cervix, 3-serosal slivers, 4-5-uterine wall, 6-endomyometrium, 7- fallopian tube
- C. In formalin labeled "left fallopian tube and ovary" is a 6 x 0.6 cm fimbriated fallopian tube surfaced by congested serosa. The tube is adhesed to the attached, 4 x 3 x 2 cm ovary that is remarkable for a embedded 1 cm metal clip, externally. Sectioning reveals 2, 1.5 and 2 cm cysts, that are devoid of contents. The larger cyst has a pink, congested lining. No solid masses or excrescences are identified. Sampled: 1-4-left tube and ovary
- D. In formalin labeled "left periureteral":  $0.7 \times 0.5 \times 0.2$  cm tissue piece(s). Totally submitted, 1 block.
- E. In formalin labeled "right cul-de-sac": 1,  $2 \times 0.9 \times 0.3$  cm tissue piece(s). The specimen is bisected. Totally submitted, 1 block. JRW

# Microscopic Description

A through E. Gross description reviewed and microscopic examined.