

Histology Cytology Report

Clinical Information

Preoperative diagnosis: Interstitial cystitis, Pelvic and perineal pain

Clinical data:

Postoperative diagnosis: Pending

Special instructions:

Specimen source: A. Endocervical curettings; B. Endometrial curettings; C. Left fallopian tube cyst;
D. Right fallopian tube cyst

Diagnosis

A. ENDOCERVIX, CURETTINGS:

ENDOCERVICAL GLANDULAR MUCOSA WITH NO SIGNIFICANT HISTOPATHOLOGIC ABNORMALITY.

B. ENDOMETRIUM, CURETTINGS:

PROLIFERATIVE PHASE ENDOMETRIUM.

C. LEFT FALLOPIAN TUBE CYST, EXCISION:

BENIGN SEROUS PARATUBAL CYST.

D. RIGHT FALLOPIAN TUBE CYST, EXCISION:

BENIGN SEROUS PARATUBAL CYST.

INCIDENTAL BENIGN ADRENOCORTICAL REST.

Gross Description

A. In formalin labeled "endocervical curetting" is a 1 x 0.5 x 0.2 cm aggregate of tissue fragments admixed with mucus. Totally submitted, 1 block.

B. In formalin labeled "endometrial sample" is a 3 x 1.5 x 0.2 cm aggregate of tissue fragments admixed with blood. Totally submitted, 1 block.

C. In formalin labeled "left fallopian tube cyst": 1, 0.5 cm cystic tissue piece(s). Totally submitted, 1 block.

Histology Cytology Report

Gross Description

D. In formalin labeled "right fallopian tube cyst": 2, 0.5-1.1 cm tissue piece(s). Totally submitted, 1 block. [REDACTED]

Microscopic Description

A, B, C, and D. Gross description reviewed and microscopic examined.