

Surgical Pathology Report

SPECIMEN(S) RECEIVED

A:Left ovarian adhesion
B:Left fallopian tube
C:Left ovarian fossa, biopsy
D:Left periureteral, biopsy
E:Right fallopian tube
F:Uterus

CLINICAL INFORMATION

N/A.

PRE OP DIAGNOSIS

DEEP ENDOMETRIOSIS OF THE PELVIC PERITONEUM OTHER SPECIFIED SITES PELVIC PAIN IRREGULAR PERIODS
OTHE

FINAL DIAGNOSIS

- A. Ovarian adhesion, left, biopsy:
 - Benign fibrovascular tissue
- B. Fallopian tube, left, salpingectomy:
 - Unremarkable fallopian tube
- C. Ovarian fossa, left, biopsy:
 - Fibroadipose tissue with features suggestive of endometriosis
- D. Peri-ureter, left, biopsy:
 - Endometriosis involving fibroadipose tissue
- E. Fallopian tube, right, salpingectomy:
 - Unremarkable fallopian tube
- F. Uterus, supra-cervical hysterectomy:
 - Secretory endometrium
 - Adenomyosis

GROSS DESCRIPTION

A. Specimen is labeled "left ovarian adhesion" and consists of a $0.6 \times 0.2 \times 0.1$ cm string-like, tan-white membranous tissue fragment, entirely submitted in 1 cassette.

B. Specimen is labeled "left fallopian tube" and consists of a 4.8 cm in length purple-red fallopian tube that averages 0.6 cm in diameter. The tube is partially covered by adhesed adipose tissue and has unremarkable cut surfaces. The fimbriated end and representative cross-sections are submitted in 1 cassette.

C. Specimen is labeled "left ovarian fossa" and consists of a $1.3 \times 0.4 \times 0.2$ cm sheetlike tan membranous soft tissue fragment, serially sectioned and entirely submitted in 1 cassette.

D. Specimen is labeled "left periureteral" and consists of a $1.3 \times 0.6 \times 0.2$ cm pink-tan membranous tissue fragment focally covered by yellow adipose tissue. The fragment is sectioned and entirely submitted in 1 cassette.

E. Specimen is labeled "right fallopian tube" and consists of a 5.3 cm in length purple-pink fallopian tube that averages 0.5 cm in diameter. Cut surfaces are unremarkable. The fimbriated end and representative cross-sections are submitted in 1 cassette.

F. Specimen is labeled "uterus" and consists of a uterus without cervix that is 38 g and measures 5 cm fundus to cervical resection site, 4 cm cornu to cornu and 3.5 cm anterior to posterior. The serosal surface is smooth. The endometrial cavity is 3 cm long by up to 1.5 cm wide and is lined by tan-red endometrium that averages 0.3 cm thick. The myometrium is mildly trabeculated and averages 1.5 cm thick. Intramural nodules are not identified. Representative sections are submitted in 4 cassettes as follows:

- 1: Cervical resection site
- 2-3: Anterior full-thickness
- 4: Posterior full-thickness